

Pure and Applied Mathematics Quarterly
 Volume 1, Number 4
 (Special Issue: In Memory of
 Armand Borel, Part 3 of 3)
 701—735, 2005

Average Values of Modular L -series Via the Relative Trace Formula

Dinakar Ramakrishnan and Jonathan Rogawski

In memory of Armand Borel

Contents

1. Introduction

For any pair of positive integers (N, k) with k even, let $\mathcal{F}_N(k)$ denote an orthogonal basis of holomorphic Hecke-eigencuspforms φ of weight k for the congruence subgroup $\Gamma_0(N)$, acting on the upper half plane $\mathcal{H}$. Let $\mathcal{F}_N(k)^{\text{new}}$ be the subspace of newforms. Denote by $\langle \varphi, \varphi \rangle$ the Petersson norm $\int_{\Gamma_0(N) \backslash \mathcal{H}} |\varphi(z)|^2 y^{k-2} dx dy$. Normalize the L -function of φ so that the functional equation relates s to $1 - s$.

For any prime p define probability measures $\mu_+(x)dx$ and $\mu_-(x)dx$ on the interval $[-2, 2] \subset \mathbb{R}$ by

$$\mu_+(x) = \frac{(p-1)}{2\pi} \frac{\sqrt{4-x^2}}{(p^{1/2} + p^{-1/2} - x)^2}$$

and

$$\mu_-(x) = \frac{(p+1)}{2\pi} \frac{\sqrt{4-x^2}}{((p^{1/2} + p^{-1/2})^2 - x^2)}$$

Note that μ_- is the familiar Plancherel measure at p , but (the more interesting) μ_+ is neither that nor the Sato-Tate measure. In either case, the limit of $\mu_{\pm}$ exists as $p \rightarrow \infty$ and equals the Sato-Tate measure $\frac{\sqrt{4-x^2}}{2\pi} dx$.

Now fix a primitive quadratic character χ of conductor $D < 0$, and restrict attention to the set of N which are prime, do not divide D , and satisfy $\chi(-N) = 1$. From here on assume that p does not divide DN , and let $\mu_p(x)$ be $\mu_+(x)$ or $\mu_-(x)$ depending on whether $\chi(p)$ is 1 or -1 .

Our main result is the following

Theorem A *Let $\chi = \chi_D$ be a quadratic character of conductor $D < 0$, N a prime with $\chi(-N) = 1$, k an even integer > 2 , and p an auxiliary prime $\nmid ND$ with $\chi(p) = \pm 1$. Then for any subinterval J of $[-2, 2]$, we have ($\forall \varepsilon > 0$)*

$$\sum_{\substack{\varphi \in \mathcal{F}_N(k)^{\text{new}} \\ a_p(\varphi) \in J}} \frac{1}{(\varphi, \varphi)} L\left(\frac{1}{2}, \varphi \otimes \chi\right) L\left(\frac{1}{2}, \varphi\right) = 2\mu_p(J)c_k L(1, \chi) + O(N^{-k/2+\varepsilon}),$$

where $L(s, \chi_D)$ is the Dirichlet L -function $\sum_{n \geq 1} \chi_D(n)n^{-s}$, and

$$c_k = d(\mathcal{D}_k) 2^k \pi \frac{k((k/2 - 1)!)^2}{(k - 1)!} \left(1 + \sum_{n=0}^{\frac{k}{2}-2} \binom{k}{2n+1} (-1)^{k/2-n} (k/2 + n - 1)! (k/2 - n - 2)! \right),$$

with $d(\mathcal{D}_k)$ denoting the formal degree of the holomorphic discrete series representation of $\mathrm{PGL}(2, \mathbb{R})$ of weight k .

Here $a_p(\varphi)$ denotes the (normalized) p -Hecke eigenvalue of φ , which is known by Deligne to lie in $[-2, 2]$. If c_p is the usual p -Hecke eigenvalue, which is an algebraic integer, a_p equals $c_p/p^{(k-1)/2}$. Concerning the formal degree, when the Haar measure $dg = dg_\infty \otimes dg_0$ on $\mathrm{GL}(2, \mathbb{A})$ is chosen so that dg_∞ corresponds to the Poincaré measure $dx dy/y^2$ on the upper half plane, $\mathcal{D}_k = (k - 1)/2$.

When J is the full interval $[-2, 2]$, there is no issue with the measure, and such a result has then been known for some time (for $k = 2$) by the work of W. Duke ([Du]); see also [Lu]. Our work began with our effort to understand these papers. It should be mentioned that in a recent paper of Iwaniec and Sarnak ([IwS]), they introduce a novel program to prove a good lower bound for $L(1, \chi)$ from a simultaneous lower bound, for a family of φ , of $L(\frac{1}{2}, \varphi)$ and $L(\frac{1}{2}, \varphi \otimes \chi)$. There are other notable papers dealing with non-vanishing results for modular L -functions and their derivatives; see for example [KMV].

Our approach here is representation-theoretic, and makes use of the *relative trace formula* of H. Jacquet; more precisely, we study the integral of the $\mathrm{GL}(2)$ kernel over the square of the maximal split torus T . In the process we are led to compute and interpret some period integrals (and certain related local ones) explicitly, and this is where the technical difficulty resides. For simplicity of exposition, we are not treating the case $k = 2$ here. The reason is that in our method we need to work with a special, factorizable function f on $\mathrm{GL}(2, \mathbb{A})$ with f_∞ being the matrix coefficient of the discrete series of weight k , which is

unfortunately *not* integrable for $k = 2$. For $k > 2$, this choice of f_∞ in the relative trace formula allows us to isolate the cusp forms of weight k (and level N), and for $k = 2$, we will need to modify f_∞ à la Hecke, which will force us to consider other cusp forms and Eisenstein series which have the corresponding K_∞ -type. After subtracting off a suitable singularity on either side of the trace formula, we will be left with a similar asymptotic formula for $k = 2$ as well.

Here is a concrete consequence of our theorem:

Corollary B *Fix p, χ as above, with D being the conductor of χ , and an even integer $k > 2$. Choose any non-empty interval $J \in [-2, 2]$. Then there are infinitely many primes N not dividing pD , which are inert in $K = \mathbb{Q}[\sqrt{D}]$, such that for a holomorphic newform φ of level N and weight k ,*

- (i) $a_p \in J$; and
- (ii) $L(1/2, \varphi \otimes \chi)L(1/2, \varphi) \neq 0$.

In particular, there are cusp forms φ of weight k and prime level for which a_p lies arbitrarily close to 2 or -2 , but at the same time having $L(1/2, \varphi \otimes \chi)L(1/2, \varphi)$ stay non-zero. Without the requirement on the non-vanishing of the L -value, this is a well known result of Serre ([Se]); see also Sarnak ([Sa]) in the context of Maass forms.

Finally, we note that, unlike in the Sato-Tate and the Plancherel cases, there is a *bias* built into our measure μ_+ . It favors positive x .

Much of the work on this paper was done five years ago, but the explicit nature of the measure was worked out only recently. One can extend the result without much trouble to square-free level N with $\chi(-N) = 1$, and also to analogous situations over totally real fields, even with varying weights ≥ 2 at the different infinite places, but of the same parity. (If $\chi(-N) = -1$, the relevant L -function vanishes at $s = 1/2$ and in that case one should consider the derivative.) However, the general level N case is more difficult. One can similarly treat the case of Maass forms, but then the infinity type λ will not be fixed, but will lie in a short interval. We thank Nathaniel Grossman for helpful comments on the evaluation of an archimedean integral involving the hypergeometric function. The first author also thanks Bill Duke, Jeff Hoffstein, Hervé Jacquet, Wenzhi Luo, Philippe Michel, and S. Rallis for interesting conversations. After this paper was finalized, we heard from Philippe Michel of a similar work of E. Royer (Bull. Soc. Math. France 128 (2000), no. 2, 219–248) involving the average of the *single* L -function $L(1/2, \varphi)$, without using representation theory. Last but not least, we acknowledge with thanks the support from the National Science Foundation.

2. The geometric side: Isolation of the main term

2.1. Regularization and the main integral. Let G denote $\mathrm{GL}(2)/\mathbb{Q}$ with center Z , and set $\tilde{G} = G/Z$. Consider the kernel

$$K(x, y) = \sum_{\gamma \in \tilde{G}(\mathbb{Q})} f(x^{-1}\gamma y)$$

where f is a factorizable, smooth function on $\tilde{G}(\mathbb{A})$ with some good properties (see section 2.3). If T denotes the maximal split torus of G , put $\tilde{T} = T/Z$. The relative trace formula which we want to make use of, and which is due to Jacquet ([Ja]), involves integration of this kernel over the square of $\tilde{T}(\mathbb{A})/\tilde{T}(\mathbb{Q})$ against a character. Unfortunately, this integral does not make sense. In order to take care of the convergence problem, we regularize the situation.

First a few words about the normalization of Haar measures. Let $\mathbb{A}$ denote the ring of adeles of $\mathbb{Q}$, and let $I_{\mathbb{Q}} = \mathbb{A}^*$ be its multiplicative group of ideles. Recall that $\mathbb{Q}^*$ is a discrete subgroup of $\mathbb{A}^*$, with the quotient $\mathbb{A}^*/\mathbb{Q}^*$ being identified with $\mathbb{R}_+^* \times \prod_q \mathbb{Z}_q^*$, where q runs over all the primes. Choose the measure $dx/|x|$ on $\mathbb{R}^*$, with dx denoting the Haar measure on $\mathbb{R}$. Normalize the measure on each $\mathbb{Q}_q^*$ so that $\mathbb{Z}_q^*$ gets volume 1. Then if $I_{\mathbb{Q}}$ denote the kernel of norm on $I_{\mathbb{Q}}$, the compact group $I_{\mathbb{Q}}^1/\mathbb{Q}^*$ gets volume 1 under the quotient measure. Finally, we will normalize, for convenience, the measures on K_q and $Z_q = Z(\mathbb{Q}_q)$ (for every prime q) such that $K_q Z_q / Z_q$ gets volume 1.

Now let χ be a non-trivial, quadratic idele class character of $\mathbb{Q}$. For $s_1, s_2 \in \mathbb{C}$, set

$$I_c(f; s_1, s_2) = \int_c \int_c K\left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} b & 0 \\ 0 & 1 \end{pmatrix}\right) \chi(a) |a|^{s_1} |b|^{s_2} d^*a d^*b.$$

where the subscript c indicates that the integrals are taken over $a, b \in \mathbb{Q}^* \backslash \mathbb{A}^*$ such that $c^{-1} < |a|, |b| < c$.

We are interested in the following limits:

$$I(f; s_1, s_2) := \lim_{c \rightarrow \infty} I_c(f; s_1, s_2).$$

and

$$I(f) := \lim_{s_1, s_2 \rightarrow 0} I(f; s_1, s_2)$$

Let T be the diagonal subgroup. Set

$$\tilde{T} = \{t_a\} \quad \text{where} \quad t_a = \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}.$$

Clearly, $\tilde{T}$ identifies with $T/Z \subset G/Z$.

For $\delta \in \tilde{G}$, define the subgroup

$$C_\delta = \{(t, t') \in \tilde{T} \times \tilde{T} : t^{-1}\delta t' = z\delta \quad \text{for some } z \in Z\}.$$

Then

$$I_c(f; s_1, s_2) = \sum_{\{\delta\}} I_c(\delta, f; s_1, s_2)$$

where $\{\delta\}$ is a set of representatives for the double cosets $\tilde{T} \backslash \tilde{G} / \tilde{T}$ and

$$I_c(\delta, f; s_1, s_2) = \int_{C_\delta(\mathbb{Q}) \backslash (\tilde{T}(\mathbf{A}) \times \tilde{T}(\mathbf{A}))}^c f(t_a^{-1} \delta t_b) \chi(a) |a|^{s_1} |b|^{s_2} d^*a d^*b$$

where the superscript c indicates that the integral is taken over $c^{-1} < |a|, |b| < c$.

2.2. Coset representatives and centralizers. Define matrices

$$\xi(x) = \begin{pmatrix} 1 & x \\ 1 & 1 \end{pmatrix}$$

and

$$n^+ = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad n^- = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \quad \varepsilon = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

Further let e denote the identity matrix in G .

Observe that $T \backslash G / T = \tilde{T} \backslash \tilde{G} / \tilde{T}$.

Lemma 1. *The set of matrices*

$$\{\xi(x) : x \neq 0, 1\} \cup \{e, \varepsilon, n^+, \varepsilon n^+, n^-, \varepsilon n^-\}$$

is a set of representatives for the double cosets $T \backslash G / T$.

The elements $\xi(x)$ and the orbits they represent will be called *regular*. The six remaining representatives and their orbits will be called *singular*.

We have

$$\begin{pmatrix} a^{-1} & \\ & 1 \end{pmatrix} \begin{pmatrix} x & y \\ z & w \end{pmatrix} \begin{pmatrix} b & \\ & 1 \end{pmatrix} = \begin{pmatrix} ba^{-1}x & a^{-1}y \\ bz & w \end{pmatrix}.$$

Lemma 2. $C_\delta = \{e\}$ if δ is regular or if $\delta \in \{n^+, \varepsilon n^+, n^-, \varepsilon n^-\}$. On the other hand,

$$C_e = \{(t_a, t_a) : a \in I_{\mathbb{Q}}\}$$

$$C_\varepsilon = \{(t_a, t_a^{-1}) : a \in I_{\mathbb{Q}}\}.$$

The proof is straight-forward and will be left to the reader.

Lemma 3. *As χ is non-trivial, we have $(\forall c > 0)$*

$$I_c(e, f) = I_c(\varepsilon, f) = 0.$$

Proof. To treat $\delta = e$, write a typical element of $\tilde{T} \times \tilde{T}$ as $(t_x t_a, t_a)$. Then we may write $I_c(e, f)$ as a double integral

$$\int_{\mathbb{Q}^* \setminus \mathbb{A}^*, c^{-1} < |a| < c} \left(\int_{x \in \mathbb{A}^*, c^{-1} < xa < c} f(t_x^{-1}) \chi(x) |x|^{s_1} d^* x \right) \chi(a) |a|^{s_1 + s_2} d^* a.$$

Since χ is non-trivial and of finite order, its restriction to $I_{\mathbb{Q}}^1$ is also non-trivial. Consequently its integral over $a \in \mathbb{Q}^* \setminus I_{\mathbb{Q}}^1$ vanishes, showing that $I_c(e, f) = 0$.

To treat $\delta = \varepsilon$, write a typical element of $\tilde{T} \times \tilde{T}$ as $(t_x t_a, t_a^{-1})$. Then we may write $I_c(\varepsilon, f)$ as a double integral

$$\int_{\mathbb{Q}^* \setminus \mathbb{A}^*, c < |a| < c^{-1}} \left(\int_{x \in \mathbb{A}^*, c^{-1} < xa < c} f(t_x^{-1}) \chi(x) |x|^{s_1} d^* x \right) \chi(a) |a|^{s_1 - s_2} d^* a.$$

Again, the integral of χ over $a \in \mathbb{Q}^* \setminus I_{\mathbb{Q}}^1$ vanishes, implying that $I_c(\varepsilon, f) = 0$. □

2.3. The test function. Let S' be the set of finite primes q at which χ is ramified. Fix two distinct primes $p, N \notin S'$ and set

$$S = S' \cup \{p, N, \infty\}.$$

We take our function to be of the form

$$f = f_{\infty} \times f_p \times f_N \times f_{S'} \times f^S$$

where $f_{S'} = \prod_{v \in S'} f_v$ and $f^S = \prod_{v \notin S} f_v$.

Let $\mathcal{D}_k$ be the holomorphic discrete series representation of $\mathrm{PGL}(2, \mathbb{R})$ of lowest weight k . We choose the archimedean function f_{∞} to be the formal degree $d(\mathcal{D}_k)$ times the complex conjugate of the matrix coefficient

$$\langle \mathcal{D}_k(g)v, v \rangle,$$

where v is a lowest vector of weight k , which is unique up to a scalar. We will take v to be a unit vector. Explicitly we take

$$f_{\infty}(g) = d(\mathcal{D}_k) \frac{(2\sqrt{\det g})^k}{(a + d + i(b - c))^k} \quad \text{if} \quad g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad \text{and} \quad \det g > 0$$

and $f_\infty(g) = 0$ if $\det g < 0$. This function takes the value $d(\mathcal{D}_k)$ at the identity matrix and has the right K_∞ -type, as needed.

We take the function f_p to be an arbitrary element in the bi- $Z_p K_p$ -invariant Hecke algebra of $G(\mathbb{Q}_p)$. The function f_N is the characteristic function of the subgroup $Z_N K_0(N)$ divided by the measure V_N of $Z_N K_0(N)/Z_N$, where

$$K_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2(\mathbf{Z}_N) : c \equiv 0 \pmod{N} \right\}.$$

We will choose the function f^S to be the characteristic function of $\prod_{v \notin S} Z_v GL_2(\mathbf{Z}_v)$.

Finally let $v = q$ be in S' . If q^m is the conductor of χ_q , denote by X the set of representatives in $q^{-m}\mathbb{Z}_q \subset \mathbb{Q}_q$ for $q^{-m}\mathbb{Z}_q/\mathbb{Z}_q$, which is a finite group isomorphic to $\mathbb{Z}/q^m$. We may view χ_v as a character of X . Put

$$f_v = g(\chi_v)^{-1} \sum_{z \in X} \chi_{1,v}(z) f_{z,v},$$

with $f_{z,v}$ being the characteristic function of $\begin{pmatrix} 1 & z \\ 0 & 1 \end{pmatrix} K_v Z_v$, and $g(\chi_v)$ the *Gauss sum*

$$g(\chi_v) = \int_{\mathbb{Z}_q^*} \chi_v(x) \psi_v(q^{-m}x) d^*x.$$

Here ψ denotes the additive character of $\mathbb{Q}_q$ defined as the composite

$$\mathbb{Q}_q \rightarrow \mathbb{Q}_q/\mathbb{Z}_q \rightarrow \mathbb{Q}/\mathbb{Z} \rightarrow S^1,$$

with the last arrow on the right being $x \rightarrow e^{2\pi i x}$. It is well known that $g(\chi_v)$ has absolute value $q^{m/2}$. Similarly, the global Gauss sum $g(\chi)$, which is a product of local ones, has absolute value $|D|^{1/2}$ since $|D|$ is the conductor of χ . In fact we have, due to the oddness of χ ,

$$g(\chi) = i|D|^{1/2},$$

which is the unique square-root of D in $\mathbb{C}$ with positive imaginary part.

The integral $I(\delta, f)$ factors as a product:

$$I(\delta, f) = I_\infty(\delta, f) I_p(\delta, f) I_N(\delta, f) I_{S'}(\delta, f) I^S(\delta, f),$$

where $I^S(\delta, f)$ (resp. $I_{S'}(\delta, f)$) is the product of $I_v(\delta, f)$ over places $v \notin S$ (resp. $v \in S'$).

2.4. The dominant terms. It will turn out that since χ is non-trivial, the terms corresponding to $\delta = n^\pm$ will be the dominant ones when the level becomes large. We will first evaluate $I(n^+, f)$, which is a limit of $I(n^+, f; s_1, s_2)$ as $s_1, s_2 \rightarrow 0$.

Theorem 2.1. *We have*

$$I(n^+, f) = \frac{\pi k}{2^k d(\mathcal{D}_k) \pi V_N} \left(\sum_{n=0}^{k/2-2} \binom{k+2n-1}{2n+1} \right) F_p(\chi) L(1, \chi),$$

where

$$F_p(\chi) := \lim_{s_1, s_2 \rightarrow 0} \frac{I_p(n^+, f; s_1, s_2)}{L(s, \chi_p)}$$

and

$$V_N := \text{vol}(Z_N K_0(N)/Z_N)$$

Note that $L(s, \chi_p)$ has a simple pole at $s = 0$ iff $\chi(p) = 1$.

For any place v , the local integral at v can be written as:

$$\begin{aligned} I_v(n^+; s_1, s_2) &= \iint_{\mathbb{Q}_v^* \times \mathbb{Q}_v^*} f_v\left(\begin{pmatrix} ab & a \\ 0 & 1 \end{pmatrix}\right) \chi_v(a)^{-1} |a|^{-s_1} |b|^{s_2} d^*a d^*b \\ &= \iint_{\mathbb{Q}_v^* \times \mathbb{Q}_v^*} f_v\left(\begin{pmatrix} b & a \\ 0 & 1 \end{pmatrix}\right) \chi_v(a) |a|^{-s_1-s_2} |b|^{s_2} d^*a d^*b. \end{aligned}$$

Theorem 2.1 is a consequence of the following, thanks to the functional equation of $L(s, \chi)$, which implies $L(0, \chi) = \frac{|D|^{1/2}}{\pi} L(1, \chi)$, and the fact that $|D|$ is the global conductor of χ :

Proposition 2.1. **(a):** *Let v be a place $\notin \{p, N, \infty\}$. If χ_v is unramified (resp. ramified of conductor q_v^m), we have:*

$$I_v(n^+; s_1, s_2) = L_v(-s_1 - s_2, \chi_v),$$

(resp. $I_v(n^+; s_1, s_2) = g(\chi_v)^{-1} L_v(-s_1 - s_2, \chi_v)$), implying that

$$I(n^+, f) = g(\chi)^{-1} F_\infty(\chi) F_N(\chi) F_p(\chi) L(0, \chi),$$

where for $v = \infty, p, N$,

$$F_v(\chi) = \lim_{s_1, s_2 \rightarrow 0} \frac{I_v(n^+, f; s_1, s_2)}{L(-s_1 - s_2, \chi_v)}$$

(b): *At $v = N$, we have*

$$I_v(n^+; s_1, s_2) = \frac{1}{V_N} L_N(-s_1 - s_2, \chi_N);$$

(c): At the archimedean place, since $\chi_\infty = \text{sgn}$, the local factor $L(s, \chi_\infty) = \pi^{-(s+1)/2} \Gamma(\frac{s+1}{2})$ is regular at $s = 0$ with value 1, and we have

$$F_\infty(n^+) = I_\infty(n^+) = -\frac{2^k i \pi k ((k/2 - 1)!)^2 h(k)}{(k-1)!},$$

where

$$h(k) := 1 + \sum_{n=0}^{\frac{k}{2}-2} \binom{k}{2n+1} (-1)^{k/2-n} (k/2 + n - 1)! (k/2 - n - 2)!$$

Remark: Note that $F_\infty(n^+)$ is purely imaginary, and so is the Gauss sum $g(\chi)$, because χ is odd. It follows that the global geometric term $I(n^+)$ is real. It is not hard to see that it is in fact positive.

Proof. (a) First let $v = q$ be outside $S = S' \cup \{\ell, N, \infty\}$, corresponding to a prime $q \neq \ell, N$. By our choice of f , f_v is the characteristic function of $Z_v K_v$. Consequently,

$$(a, b) \rightarrow f_v \begin{pmatrix} b & a \\ 0 & 1 \end{pmatrix}$$

is simply the characteristic function of $\mathbf{Z}_v \times \mathbf{Z}_v^*$. Also, χ is unramified at v . So we get

$$I_v(n^+) = \sum_{n \geq 0} \chi(q^{-n}) q^{n(s_1+s_2)}.$$

This is a geometric series summing to $(1 - \chi(q)^{-1} q^{s_1+s_2})^{-1}$, whence the assertion of part (a) in this case.

Next suppose $v = q$ is in S' , with q^m being the conductor of χ_v . By construction,

$$I_v(n^+) = g(\chi_v)^{-1} \sum_{z \in X} I_{z,v}(n^+),$$

where

$$I_{z,v}(n^+) = \chi_v(z) \iint_{\mathbb{Q}_v^* \times \mathbb{Q}_v^*} f_{z,v} \begin{pmatrix} b & a \\ 0 & 1 \end{pmatrix} \chi_v(a^{-1}) |a|^{-s_1-s_2} |b|^{s_2} d^* a d^* b.$$

Only those z with invertible images in $q^{-m} \mathbb{Z}_q / \mathbb{Z}_q$ give a non-zero contribution. So we will restrict our attention to these.

Recall that f_v has support on $\begin{pmatrix} 1 & z \\ 0 & 1 \end{pmatrix} K_v Z_v$, and note that

$$\lambda \begin{pmatrix} 1-z & \\ 0 & 1 \end{pmatrix} \begin{pmatrix} b & a \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \lambda b & \lambda(a-z) \\ 0 & \lambda \end{pmatrix}.$$

Then it follows that for $f_{z,v} \begin{pmatrix} b & a \\ 0 & 1 \end{pmatrix}$ to not vanish, we need $b \in \mathbb{Z}_q^*$ and $a - z \in \mathbb{Z}_q$.

Thus a lies in $q^{-m}\mathbb{Z}_q$ and has the same image as z in $q^{-m}\mathbb{Z}_q/\mathbb{Z}_q$. Consequently, since χ_v has conductor q^m and thus the pullback of a character of $(q^{-m}\mathbb{Z}_q/\mathbb{Z}_q)^* \simeq (\mathbb{Z}/q^m)^*$, we must have $\chi_v(z)\chi_v(a^{-1}) = 1$. We get

$$I_{z,v}(n^+) = \text{vol}(1 + q^m\mathbb{Z}_q),$$

when z has invertible image in $\mathbb{Z}/q^m$. The assertion follows once we note:

- (i) $\varphi(q^m)\text{vol}(1 + q^m\mathbb{Z}_q)$ equals $\text{vol}(\mathbb{Z}_q^*) = 1$, and
- (ii) $L(s, \chi_v) = 1$ as χ_v is ramified.

(b) Let $v = N$. Recall that f_N is the characteristic function of $Z_N K_0(N)$ divided by V_N . Like in (a), the function

$$(a, b) \rightarrow f_N \begin{pmatrix} b & a \\ 0 & 1 \end{pmatrix}$$

is the characteristic function of $\mathbf{Z}_N \times \mathbf{Z}_N^*$, but divided by V_N . The assertion follows. (When we consider n^- , the situation will be slightly different.)

(c) Let $v = \infty$. Recall that $\chi_\infty(-1) = -1$. By the definition of f_∞ , its value on $\begin{pmatrix} b & a \\ 0 & 1 \end{pmatrix}$ is $d(\mathcal{D}_k) \frac{b^{k/2}}{(b+1-ia)^k}$ (resp. 0) when b is positive (resp. negative). Noting that $d^*x = dx/|x|$ on $\mathbb{R}^*$. We get

$$\begin{aligned} d(\mathcal{D}_k)^{-1} I_\infty(n^+; s_1, s_2) &= 2^k \int_{-\infty}^{\infty} \int_0^{\infty} \frac{b^{k/2+s_2-1} |a|^{-s_1-s_2-1} \text{sgn}(a)}{(b+1-ia)^k} dadb \\ &= 2^k \int_{-\infty}^{\infty} \int_0^{\infty} \frac{b^{k/2+s_2-1} |a|^{-s_1-s_2-1} \text{sgn}(a) ((b+1)+ia)^k}{(a^2 + (b+1)^2)^k} dadb, \end{aligned}$$

where b runs over $(0, \infty)$ and a runs over $(-\infty, \infty)$. Appealing to the binomial expansion

$$((b+1)+ia)^k = \sum_{j=0}^k \binom{k}{j} i^{k-j} a^{k-j} (b+1)^j,$$

we may write

$$d(\mathcal{D}_k)^{-1} I_\infty(n^+; s_1, s_2) = 2^k \sum_{j=0}^k \binom{k}{j} i^{k-j} I_j,$$

where

$$I_j = \int_{-\infty}^{\infty} \int_0^{\infty} \frac{b^{k/2+s_2-1} |a|^{k-j-s_1-s_2-1} (a/|a|)^{1+k-j} (b+1)^j}{(a^2 + (b+1)^2)^k} dadb.$$

Note that that if j , and hence $k - j$, is even, then the integrand is odd in the a -variable and so I_j vanishes. So we may, and we will, assume from hereon that j is odd. We have

$$\begin{aligned} I_j &= 2 \int_0^\infty \int_0^\infty \frac{b^{k/2+s_2-1} a^{k-j-s_1-s_2-1} (b+1)^j}{(a^2 + (b+1)^2)^k} da db \\ &= 2 \int_0^\infty \int_0^\infty \frac{b^{k/2+s_2-1} a^{k-j-s_1-s_2-1} (b+1)^{j-2k}}{((a/(b+1))^2 + 1)^k} da db \\ &= 2 \int_0^\infty \int_0^\infty \frac{b^{k/2+s_2-1} a^{k-j-s_1-s_2-1} (b+1)^{-k-s_1-s_2}}{(a^2 + 1)^k} da db. \end{aligned}$$

So the integral factors as

$$(0) \quad I_j = I_{j,1} I_{j,2},$$

where

$$I_{j,1} = 2 \int_0^\infty \frac{a^{k-j-s_1-s_2-1}}{(a^2 + 1)^k} da$$

and

$$I_{j,2} = \int_0^\infty b^{k/2+s_2-1} (b+1)^{-k-s_1-s_2} db.$$

Note that $I_{j,2}$ is in fact independent of j .

By the substitution $u = a^2$, we get

$$I_{j,1} = \int_0^\infty \frac{u^{(k-j-s_1-s_2)/2-1}}{(u+1)^k} du.$$

Recall that the Beta function $B(z, w) = \frac{\Gamma(z)\Gamma(w)}{\Gamma(z+w)}$ has the following integral representations (cf. [A-S], p.258):

$$(*) \quad B(z, w) = \int_0^\infty t^{z-1} (1+t)^{-z-w} dt = \int_0^\infty \frac{t^{z-1}}{(1+t)^{z+w}} dt.$$

It follows immediately that

$$(1) \quad I_{j,1} = B((k-j-s_1-s_2)/2, (k+j+s_1+s_2)/2)$$

Now let $v = b/(b+1)$ in the expression for $I_{j,2}$, so that $b = v/(1-v)$ and $db = dv/(1-v)^2$. We obtain:

$$I_{j,2} = \int_0^1 v^{k/2-1+s_2} (1-v)^{k/2+s_1-1} dv.$$

Applying the first identity of (*) we get

$$(2) \quad I_{j,2} = B(k/2 + s_2, k/2 + s_1).$$

Putting (0), (1) and (2) together, and writing everything in terms of the Γ -function, we finally obtain

$$(3) \quad I_j = \frac{\Gamma((k-j-s_1-s_2)/2)\Gamma((k+j+s_1+s_2)/2)\Gamma(k/2+s_2)\Gamma(k/2+s_1)}{\Gamma(k)\Gamma(k+s_1+s_2)}.$$

Consequently, we see that $d(\mathcal{D}_k)^{-1}I_\infty(n^+; s_1, s_2)$ equals

$$2^k \left(\sum_{\substack{j=0 \\ j \equiv 1(2)}}^k \binom{k}{j} i^{k-j} \Gamma\left(\frac{k-j-s_1-s_2}{2}\right) \Gamma\left(\frac{k+j+s_1+s_2}{2}\right) \right) \frac{\Gamma(\frac{k}{2}+s_1)\Gamma(\frac{k}{2}+s_2)}{\Gamma(k)\Gamma(k+s_1+s_2)}.$$

Now observe that for any positive integer r ,

$$\Gamma\left(\frac{1+2r}{2}\right) = (r-1)!\Gamma\left(\frac{1}{2}\right).$$

Since $\Gamma(\frac{1}{2}) = \sqrt{\pi}$ and $\Gamma(r) = (r-1)!$, putting $k = 2m, j = 2n+1$ (so that $i^{k-j} = -i(-1)^{m-n}$), we get (by sending s_1, s_2 to 0):

$$I_\infty(n^+) = -2^k i \pi d(\mathcal{D}_k) \frac{(2m)((m-1)!)^2}{(2m-1)!} \left(1 + \sum_{n=0}^{m-2} \binom{2m}{2n+1} (-1)^{m-n} (m+n-1)!(m-n-2)! \right)$$

This proves (c). □

Next we consider the case $\delta = n^-$. In this case, the local integral at any v equals

$$\begin{aligned} I_v(n^-; s_1, s_2) &= \iint_{F_v^* \times F_v^*} f_v\left(\begin{pmatrix} ab & 0 \\ b & 1 \end{pmatrix}\right) \chi_v(a)^{-1} |a|^{-s_1} |b|^{s_2} d^*a d^*b \\ &= \iint_{\mathbb{Q}_v^* \times \mathbb{Q}_v^*} f_v\left(\begin{pmatrix} a & 0 \\ b & 1 \end{pmatrix}\right) \chi_v(a^{-1}b) |a|^{-s_1} |b|^{s_1+s_2} d^*a d^*b. \end{aligned}$$

Proposition 2.2. (a): If v is a place outside $\{p, N, \infty\}$, we have:

$$I_v(n^-; s_1, s_2) = I_v(n^+; -s_2, -s_1);$$

(b): At $v = N$, we have

$$I_N(n^-; s_1, s_2) = \frac{1}{V_N} \chi(N) N^{-s_1-s_2} L(s_1+s_2, \chi_N);$$

(c): At the archimedean place we have:

$$I_\infty(n^-) = -I_\infty(n^+)$$

Since $\chi(N) = -1$, the sign change at infinity is compensated for by the sign change at N , when we go from n^+ to n^- . We get the following

Corollary 1. (A)

$$I(n^-, f)F_p(\chi) = I(n^+, f)F_p^-(\chi),$$

where

$$F_p^-(\chi) := \lim_{s_1, s_2 \rightarrow 0} \frac{I_p(n^-, f; s_1, s_2)}{L(s, \chi_p)}$$

(B) When f_p is the characteristic function of $Z_p K_p$, we have

$$F_p(\chi) = F_p^-(\chi) = 1.$$

Proof of Proposition. Part (a) is immediate. At N , the function F_N when restricted to

$$(a, b) \rightarrow f_N \begin{pmatrix} a & 0 \\ b & 1 \end{pmatrix}$$

is simply the characteristic function of $\mathbb{Z}_N^* \times N\mathbb{Z}_N$ divided by V_N . Since χ is unramified at N , $\chi(a^{-1}b)$ is just $\chi(b)$. Now part (b) follows by switching to the variable b' with $b = Nb'$. At ∞ , we obtain

$$I_\infty(n^-; s_1, s_2) = d(\mathcal{D}_k)2^k \iint_{\mathbb{R}^* \times \mathbb{R}^*} f_\infty \begin{pmatrix} a & 0 \\ b & 1 \end{pmatrix} \operatorname{sgn}(a^{-1}b) |a|^{-s_1} |b|^{s_1+s_2} d^* a d^* b,$$

and by the definition of f_∞ , it is zero if $a < 0$ and for $a > 0$ it takes the value $d(\mathcal{D}_k) \frac{a^{k/2}}{((a+1)+ib)^k}$. Comparing with what we began with at ∞ for n^+ , we see immediately (by reversing the roles of a and b) that

$$I_\infty(n^-; s_1, s_2) = -I_\infty(n^+; -s_2, -s_1).$$

The assertion of part (c) follows by sending s_1, s_2 to 0. □

2.5. The remaining singular terms. We have already seen that under our hypotheses, the global integrals $I_c(e, f)$ and $I_c(\varepsilon, f)$ are zero for any $c > 0$. So

$$I(e, f) = I(\varepsilon, f) = 0.$$

So it remains only to evaluate $I(\varepsilon n^\pm, f)$.

Proposition 2.3. *Let $\delta \in \{\varepsilon n^+, \varepsilon n^-\}$. Then $I(\delta, f) = 0$.*

Proof. As $I(\delta, f) = \prod_v I_v(\delta, f)$, it suffices to show that $I_N(\varepsilon n^\pm, f) = 0$.

First consider the $\delta = \varepsilon n^+$ case.

We have

$$I_N(\varepsilon n^+) = \iint_{\mathbb{Q}_N^* \times \mathbb{Q}_N^*} f_N\left(\begin{pmatrix} 0 & a \\ b & 1 \end{pmatrix}\right) \chi(a)^{-1} |a|^{-s_1} |b|^{s_2} d^*a d^*b$$

By definition, f_N is supported on $Z_N K_0(N)$. So $f_N\left(\begin{pmatrix} 0 & a \\ b & 1 \end{pmatrix}\right)$ is non-zero for some $(a, b) \in (\mathbb{Q}_N^*)^2$ iff there is a $\lambda \in \mathbb{Q}_N^*$ such that $\lambda, \lambda a \in \mathbb{Z}_N$, $\lambda b \in N\mathbb{Z}_N$ and $\det(\lambda \begin{pmatrix} 0 & a \\ b & 1 \end{pmatrix}) = -\lambda^2 ab \in \mathbb{Z}_N^*$. These conditions cannot be satisfied simultaneously. Done.

Now let $\delta = \varepsilon n^-$. Then we have

$$I_N(\varepsilon n^-) = \int_{(\mathbb{Q}_N^*)^2} f_N\left(\begin{pmatrix} ab & a \\ b & 0 \end{pmatrix}\right) \chi(a)^{-1} |a|^{-s_1} |b|^{s_2} d^*a d^*b$$

Again this can be non-zero only if we can find, for each a, b , a λ in $\mathbb{Q}_N^*$ such that $\lambda ab, \lambda a \in \mathbb{Z}_N$, $\lambda b \in N\mathbb{Z}_N$ and $\lambda^2 ab \in \mathbb{Z}_N^*$, which is impossible. □

To recapitulate, four of the six singular terms contribute zero to the geometric side of the relative trace formula. The remaining two, which give the dominant terms, are both non-zero, and in fact, thanks to the functional equation of $L(s, \chi)$, have the *same* limiting expression, which is a non-zero multiple of $L(1, \chi)$, as s_1, s_2 both go to 0.

2.6. The regular terms. Recall that the regular double cosets are represented by the matrices $\{\xi(x)\}$, with $x \in \mathbb{P}^1(\mathbb{Q}) - \{0, 1, \infty\}$. By abuse of notation we will write $I(x)$ for $I(\xi(x))$. At any place v , we have by definition,

$$I_v(x) = \int_{\mathbb{Q}_v^* \times \mathbb{Q}_v^*} f_v\left(\begin{pmatrix} ab & ax \\ b & 1 \end{pmatrix}\right) \chi_v(a)^{-1} |a|^{-s_1} |b|^{s_2} d^*a d^*b$$

Proposition 2.4. *Let $x \in \mathbb{P}^1(\mathbb{Q}) - \{0, 1, \infty\}$.*

(a): *Let $v = q$ be a prime not in S . When $v(1 - x) > 0$, $I_v(x)$ vanishes. Suppose $v(1 - x) = 0$, resp. $v(1 - x) < 0$. Then $v(x) \geq 0$, resp. $v(x) =$*

$v(1-x)$, and we have:

$$I_v(x) = \sum_{n=0}^{v(x)} \sum_{m=-v(x)}^{-n} ' (\bar{\chi}(q)q^{s_1})^m q^{-ns_2},$$

resp.

$$I_v(x) = \sum_{n=v(x)}^0 \sum_{m=0}^{-n} ' (\bar{\chi}(q)q^{s_1})^m q^{-ns_2},$$

where the prime on the inside sum implies (in either case) that the summation is over m of the same parity as $n + v(1-x)$. In particular,

$$v(x) = v(1-x) = 0 \implies I_v(x) = \text{vol}(\mathbb{Z}_v^*)^2.$$

(b): Let $v = q$ be a prime in S' , with q^c the conductor of χ_q . Then if $v(1-x) > c$, $I_v(x, f)$ vanishes. When $v(1-x) \leq c$, $I_v(x, f)$ is bounded by $q^{c/2}(v(x) + c)^2 \text{vol}(\mathbb{Z}_v^*)^2$.

(c): Let $v = N$. Then $I_N(x)$ vanishes when $v_N(1-x) \neq 0$. When $v_N(1-x) = 0$, we must have $v_N(x) \geq 1$ and

$$I_N(x) = \sum_{n=1}^{v_N(x)} \sum_{m=-v_N(x)}^{-n} ' (\bar{\chi}(N)N^{s_1})^m N^{-ns_2}.$$

(d): Let $v = p$, and f_p the characteristic function of $ZK \begin{pmatrix} p^r & 0 \\ 0 & p^{r'} \end{pmatrix} K$ for some integers r, r' such that $r \geq r'$. Then $I_p(x, f)$ is zero unless $v(1-x) < r + r'$, in which case it is bounded by $C(f_p)v(x)^2 \text{vol}(\mathbb{Z}_v^*)^2$, for a constant $C(f_p)$.

(e): Let $v = \infty$. Assume that $x > 0$ and that $\Re s_1, \Re s_2 \in (-k/2, k/2)$. Then, as $\chi_\infty(-1) = -1$, we have for $1-x$ is positive (resp. negative):

$$I_\infty(x) = (1-x)^{k/2} \left(I_\infty(-1, 1, 1) - (-1)^k I_\infty(-1, -1, 1) \right)$$

(resp.

$$I_\infty(x) = (x-1)^{k/2} \left(I_\infty(1, -1, -1) - (-1)^k I_\infty(1, 1, -1) \right),$$

where for $\epsilon, \delta, \nu \in \{\pm 1\}$,

$$I_\infty(\epsilon, \delta, \nu) = (-\epsilon)^{\rho-\sigma-1} \delta^{\rho+k-3\sigma-1} \nu^{k-\sigma} i^{\rho-2\sigma} B(\sigma, k-\sigma) B(\rho, k-\rho) F(k-\sigma, \rho; k; 1-\epsilon \delta \nu)$$

where $F = {}_2F_1$ is the hypergeometric function.

Proof. (a) Since $v = q$ is a prime outside S , f_v is by definition the characteristic function of $Z_v K_v$.

Lemma 4. $f_v \begin{pmatrix} ab & ax \\ b & 1 \end{pmatrix}$ is non-zero iff the following conditions are simultaneously satisfied:

- (i) $v(1-x) \leq 0$;
- (ii) $v(x) \geq v(1-x)$; in particular, $v(x) = v(1-x)$ when $v(1-x) < 0$;
- (iii) $v(1-x) - v(x) \leq v(a) \leq \min\{-v(1-x), -v(b) - v(1-x)\}$;
- (iv) $\max\{v(1-x), v(a) + v(1-x)\} \leq v(b) \leq v(x) - v(1-x)$;
- (v) $v(a) + v(b) \equiv v(1-x) \pmod{2}$.

Proof of Lemma. For $f_v \begin{pmatrix} ab & ax \\ b & 1 \end{pmatrix}$ to be non-zero, it is necessary and sufficient that there exists a λ in $\mathbb{Q}_v^*$ such that $\lambda ab, \lambda ax, \lambda b, \lambda$ are in $\mathbb{Z}_v$ and $\lambda^2 ab(1-x)$ is in $\mathbb{Z}_v^*$; in other words, one must have

$$1) \quad 2v(\lambda) + v(a) + v(b) + v(1-x) = 0;$$

$$2) \quad v(\lambda) + v(a) + v(b) \geq 0;$$

$$3) \quad v(\lambda) + v(a) + v(x) \geq 0;$$

$$4) \quad v(\lambda) + v(b) \geq 0;$$

and

$$5) \quad v(\lambda) \geq 0.$$

Eliminating $v(\lambda)$ from these, we arrive at the following system of inequalities, together with the parity condition

$$v(a) + v(b) + v(1-x) \equiv 0 \pmod{2} :$$

$$6) \quad v(a) + v(x) - v(1-x) \geq 0;$$

$$7) \quad v(b) - v(1-x) \geq 0;$$

$$8) \quad v(x) - v(1-x) \geq 0;$$

$$9) \quad v(1-x) \leq 0;$$

$$10) \quad v(a) + v(1-x) \leq 0;$$

$$11) \quad v(b) + v(1-x) - v(x) \leq 0;$$

$$12) \quad v(a) + v(b) + v(1-x) \leq 0,$$

and

$$13) \quad v(b) \geq v(a) + v(1-x).$$

To explain, 6) comes from $2) + 3) - 1)$, while 7) comes from $2) + 4) - 1)$, 8) from $3) + 4) - 1)$, 9) from $1) - 2) - 5)$, 10) from $1) - 4) - 5)$, 11) from $1) - 3) - 5)$, 12) from $1) - 2 \times 5)$, and 13) from $2 \times 4) - 1)$. The assertions of the Lemma now follows easily. □

The first consequence is that $I_v(x)$ vanishes if $v(1-x)$ is positive. Now let $v(1-x)$ be zero (resp. negative). Then $v(x) \geq 0$ (resp. $v(x) = v(1-x)$) and the inequalities (iii), (iv) of the Lemma simplify to yield the conditions

$$0 \leq v(b) \leq v(x), \quad \text{and} \quad -v(x) \leq v(a) \leq -v(b)$$

(resp.

$$v(x) \leq v(b) \leq 0 \quad \text{and} \quad 1 \leq v(a) \leq -v(x).)$$

The assertion of the Proposition now follows in the case $v = q \notin S$.

(b) Here $v = q$ is a prime where χ ramifies. If the conductor of χ_v is q^c , with $c \geq 1$, f_v is, by definition, $g(\chi_v)^{-1}$ times a χ -weighted sum of the characteristic functions $f_{z,v}$ of $\begin{pmatrix} 1 & z \\ 0 & 1 \end{pmatrix} K_v Z_v$, where z has valuation $-r$ with $0 \leq r \leq m$. We need the following:

Lemma 5. $f_{z,v} \begin{pmatrix} ab & ax \\ b & 1 \end{pmatrix}$ is non-zero only if the following conditions are simultaneously satisfied:

- (i) $v(1-x) \leq r$;
- (ii) $v(x) = v(1-x)$ when $v(1-x) < 0$, and $v(x) = 0$ when $v(1-x) > 0$;
- (iii) $v(a) \leq \min\{-v(1-x), -v(b) - v(1-x)\}$, $v(a+z) + v(ax+z) - v(a) \geq v(1-x)$;
- (iv) $\max\{v(1-x) + v(a), v(1-x) - v(a+z) + v(a)\} \leq v(b) \leq v(x) - v(1-x)$;
- (v) $v(a) + v(b) \equiv v(1-x) \pmod{2}$.

Proof of Lemma. The statement for $r = 0$ is just Lemma 5. So we may assume that $0 < r \leq c$. Then

$$\begin{pmatrix} 1 & z \\ 0 & 1 \end{pmatrix} \begin{pmatrix} ab & ax \\ b & 1 \end{pmatrix} = \begin{pmatrix} ab + bz & ax + z \\ b & 1 \end{pmatrix}$$

So for $f_{r,v} \begin{pmatrix} ab & ax \\ b & 1 \end{pmatrix}$ to be non-zero, it is necessary and sufficient that there exists a λ in $\mathbb{Q}_v^*$ such that $\lambda(a+z)b, \lambda(ax+z), \lambda b, \lambda$ are in $\mathbb{Z}_v$ and $\lambda^2 ab(1-x)$ is in $\mathbb{Z}_v^*$. In other words, one must have

$$1') \quad 2v(\lambda) + v(a) + v(b) + v(1-x) = 0;$$

$$2') \quad v(\lambda) + v(a + z) + v(b) \geq 0;$$

$$3') \quad v(\lambda) + v(ax + z) \geq 0;$$

$$4') \quad v(\lambda) + v(b) \geq 0;$$

and

$$5') \quad v(\lambda) \geq 0.$$

Eliminating $v(\lambda)$ from these, we arrive at the following system of inequalities, together with the parity condition

$$v(a) + v(b) + v(1 - x) \equiv 0 \pmod{2} :$$

$$6') \quad v(a + z) + v(ax + z) - v(a) - v(1 - x) \geq 0;$$

$$7') \quad v(b) + v(a + z) - v(a) - v(1 - x) \geq 0;$$

$$8') \quad v(ax + z) - v(a) - v(1 - x) \geq 0;$$

$$9') \quad v(1 - x) \leq v(a + z) - v(a);$$

$$10') \quad v(a) + v(1 - x) \leq 0;$$

$$11') \quad v(b) + v(1 - x) - v(x) \leq 0;$$

$$12') \quad v(a) + v(b) + v(1 - x) \leq 0;$$

and

$$13') \quad v(b) \geq v(a) + v(1 - x).$$

To explain, 6') comes from $2') + 3') - 1')$, while 7') comes from $2') + 4') - 1')$, 8') from $3') + 4') - 1')$, 9') from $1') - 2') - 5')$, 10') from $1') - 4') - 5')$, 11') from $1') - 3') - 5')$, 12') from $1') - 2 \times 5')$, and 13') from $2 \times 4') - 1')$. If $v(a) \neq v(z)$, then $v(a + z) = \min\{v(a), v(z)\}$, and so $v(a + z) - v(a) \leq 0$, and by 9'), $v(1 - x) \leq 0$. On the other hand, if $v(a) = v(z) = -r$, we get by 10') that $v(1 - x) \leq r$. This gives part (i) of the Lemma. The remaining assertions follow easily. □

So the first consequence is that $I_v(x)$ vanishes if $v(1 - x)$ is greater than r . We get from the Lemma:

$$v(1 - x) - r \leq v(b) \leq v(x) - v(1 - x).$$

This is clear when $v(a) = -r$, and if not, $v(a + z) - v(a) \leq 0$, which even gives $v(1 - x) \leq v(b)$. Next observe that when $v(1 - x)$ is 0, resp. < 0 , resp. > 0 , we have $v(x) \geq 0$, resp. $v(x) = v(1 - x)$, resp. $v(x) = 0$. We obtain

$$-r \leq v(b) \leq v(x) \quad \text{and} \quad -v(x) \leq v(a) \leq 0,$$

resp.

$$v(x) - r \leq v(b) \leq 0 \quad \text{and} \quad 0 \leq v(a) \leq -v(x) \quad \text{or} \quad v(a) = -r - v(x),$$

resp.

$$-r < v(b) < 0 \quad \text{and} \quad v(a) = -r$$

The assertion of the Proposition now follows in the case $v = q \in S'$, once we recall that $|g(\chi_v)| = q^{c/2}$.

(c) Let $v = N$. Here by construction, f_N is the characteristic function of $Z_N K_0(N)$ divided by V_N , the volume of $Z_N K_0(N)/Z_N$. So for $f_N\left(\begin{pmatrix} ab & ax \\ b & 1 \end{pmatrix}\right)$ to be non-zero, it is necessary and sufficient that there exists a λ in $\mathbb{Q}_N^*$ such that $\lambda ab, \lambda ax, \lambda$ are in $\mathbb{Z}_N$, λb is in $N\mathbb{Z}_N$ and $\lambda^2 ab(1-x)$ is in $\mathbb{Z}_N^*$. The conditions 1) through 5) above (in the proof of part (a)) remain in force except for 4), which gets replaced by

$$4'') \quad v_N(\lambda) + v_N(b) \geq 1.$$

The parity condition is the same as in (a). The only change in the inequalities 6) – 12) is that 7) (resp. 8), resp. 10)) gets replaced by

$$7'') \quad v_N(b) - v_N(1-x) \geq 1;$$

$$8'') \quad v(x) - v(1-x) \geq 1;$$

and

$$11'') \quad v(a) + v(1-x) \leq -1;$$

We again get the vanishing of $I_N(x)$ when $v_N(1-x) > 0$. Moreover, when $v_N(1-x) < 0$, we are forced to have $v_N(x) = v_N(1-x)$, which contradicts 7'); thus $I_N(x)$ vanishes in this case as well. It remains only to consider when $v_N(1-x) = 0$, in which case $v_N(x) \geq 1$. The asserted expression for $I_N(x)$ follows easily.

(d) Of course, when $r = r' = 0$, f_p is just the characteristic function of $Z_p K_p$, which was treated in detail in part (a). For general r, r' with $r \geq r'$, for $f_p\left(\begin{pmatrix} ab & ax \\ b & 1 \end{pmatrix}\right)$ to be non-zero, it is necessary and sufficient that there exists a λ in $\mathbb{Q}_v^*$ such that

$$\begin{pmatrix} \lambda ab & \lambda ax \\ \lambda b & \lambda \end{pmatrix} \in K_p \begin{pmatrix} p^r & 0 \\ 0 & p^{r'} \end{pmatrix} K_p.$$

We get analogues of conditions 1) through 12) of part (a), except for replacing the zeros on the right of those equations. In 1) and 9), replace 0 by $r + r'$, while in 2) through 4), replace 0 by r . The assertions of part d) now follow easily.

(e) Let $v = \infty$. By hypothesis, χ_∞ is *sgn*, the sign character. By the definition of f_∞ , we have

$$f_\infty \begin{pmatrix} ab & ax \\ b & 1 \end{pmatrix} = \frac{(ab(1-x))^{k/2}}{(ax-b+i(ab-1))^k} \quad (\text{resp. } 0)$$

if $ab(1-x)$ is > 0 (resp. < 0).

Suppose $1-x > 0$. Then the integral is over the first and the third quadrants. Changing variables in the third quadrant and rearranging, we get

$$I_\infty(x) = (1-x)^{k/2} \left(I'_\infty(-1, 1, 1) - (-1)^k I'_\infty(-1, -1, 1) \right),$$

where for $\epsilon, \delta, \nu \in \{\pm 1\}$,

$$I'_\infty(\epsilon, \delta, \nu) = \int_0^\infty \int_0^\infty \frac{a^{k/2-s_1-1} b^{k/2+s_2-1} da db}{(ax + \epsilon b + \delta i(ab + \nu))^k}.$$

Similarly, when $1-x$ is negative, we have

$$I_\infty(x) = (x-1)^{k/2} \left(I'_\infty(1, -1, -1) - (-1)^k I'_\infty(1, 1, -1) \right).$$

Set

$$\rho = k/2 - s_1, \quad \text{and} \quad \sigma = k/2 + s_2.$$

Then the assertion of part (c) will follow once we establish the following, which was shown to us by Nathaniel Grossman.

Lemma 6. *Suppose $k > \Re \rho > 0$ and $k > \Re \sigma > 0$. Then we have*

$$I'_\infty(\epsilon, \delta, \nu) = I_\infty(\epsilon, \delta, \nu).$$

This holds in the complex x -plane with the negative x -axis cut out.

Recall the definition of $I_\infty(\epsilon, \delta, \nu)$ from the statement of part (c) of the Proposition.

Proof. Fix ϵ, δ, ν and simply write I' for $I'_\infty(\epsilon, \delta, \nu)$. We have

$$\begin{aligned} I' &= \int_0^\infty a^{\rho-1} da \int_0^\infty \frac{b^{\sigma-1} db}{((ax + \delta \nu i) + b(\delta i a + \epsilon))^k} \\ &= \int_0^\infty \frac{a^{\rho-1} da}{(\delta i a + \epsilon)^k} \int_0^\infty \frac{b^{\sigma-1} db}{\left(\left(\frac{ax + \delta \nu i}{\delta i a + \epsilon} \right) + b \right)^k} \\ &= \int_0^\infty \frac{a^{\rho-1} \left(\frac{ax + \delta \nu i}{\delta i a + \epsilon} \right)^\sigma da}{(\delta i a + \epsilon)^k \left(\frac{ax + \delta \nu i}{\delta i a + \epsilon} \right)^k} \int_L \frac{b_1^{\sigma-1} db}{(1 + b_1)^k}, \end{aligned}$$

where L denotes the half-line defined by the positive real multiples of $\frac{\delta i a + \epsilon}{ax + \delta \nu i}$.

Since $\Re \lambda > \Re \sigma > 0$ by hypothesis, the ray of integration may be rotated back to $\mathbb{R}_+$, and the inner b_1 -integral becomes

$$\int_0^\infty \frac{b_1^{\sigma-1} db_1}{(1+b_1)^k} = B(\sigma, k-\sigma).$$

Thus

$$I' = \frac{B(\sigma, k-\sigma)}{x^{k-\sigma}(\delta i)^\sigma} J,$$

where

$$J = \int_0^\infty \frac{a^{\rho-1} da}{(a + \delta \nu i/x)^{k-\sigma} (a - i\delta \epsilon)^\sigma}.$$

Since by assumption $k > \Re \rho > 0$, the path of integration defining J can be rotated to the positive imaginary axis, giving

$$\begin{aligned} J &= \int_0^{i\infty} \frac{a^{\rho-1} da}{(a + \delta \nu i/x)^{k-\sigma} (a - i\delta \epsilon)^\sigma} \\ &= \int_0^\infty \frac{(ic)^{\rho-1} d(ic)}{(ic + \delta \nu i/x)^{k-\sigma} (ic - i\delta \epsilon)^\sigma} \\ &= i^{\rho-\sigma} \int_0^\infty \frac{c^{\rho-1} dc}{(c + \delta \nu/x)^{k-\sigma} (c - \delta \epsilon)^\sigma}. \end{aligned}$$

Case (i): $\delta \epsilon = -1$.

Put $u = \frac{c}{1+c}$ so that $c = \frac{u}{1-u}$ and $dc = \frac{du}{(1-u)^2}$. We get

$$J = i^{\rho-\sigma} (\delta \nu)^{k-\sigma} x^{k-\sigma} \int_0^1 \frac{u^{\rho-1} (1-u)^{k-\rho-1} du}{(1-u(1-\delta \nu x))^{k-\sigma}}.$$

Now we appeal to the following well known integral representation for the hypergeometric function ([A-S], formula (15.3.1), page 558):

$$F(a, b; c; z) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^\infty t^{b-1} (1-t)^{c-b-1} (1-tz)^{-a} dt,$$

for $\Re c > \Re b > 0$.

The assertion follows by putting these together.

Case (ii): $\delta \epsilon = 1$.

In this case,

$$J = i^{\rho-\sigma} \int_0^\infty \frac{c^{\rho-1} dc}{(c + \delta \nu/x)^{k-\sigma} (c-1)^\sigma}.$$

We put $c = \frac{u}{u-1}$, so that $u = \frac{c}{c-1}$ and $dc = \frac{-du}{(1-u)^2}$. We get

$$J = i^{\rho-\sigma} (\delta \nu)^{k-\sigma} x^{k-\sigma} (-1)^{\rho-\sigma-1} \int_0^\infty \frac{u^{\rho-1} (1-u)^{k-\rho-1} du}{(1-u(1+\delta \nu x))^{k-\sigma}}.$$

The assertion follows by appealing once again to the integral representation of the hypergeometric function and combining like terms.

□

3. A bound for the sum of regular terms

Put

$$I_{\text{reg}}(f) := \sum_{x \in \mathbb{Q}^* - \{1\}} I(x, f).$$

Proposition 3.1. *Let $\varepsilon > 0$. Then*

$$I_{\text{reg}}(f) \leq \frac{C}{N^{k/2-\varepsilon}},$$

for a positive constant C .

Proof. Put $t = \frac{1}{1-x}$. Since $x \rightarrow t$ is an automorphism of $\mathbb{Q}^* - \{1\}$,

$$I_{\text{reg}}(f) := \sum_{t \in \mathbb{Q}^* - \{1\}} I(x, f),$$

with $x = \frac{t-1}{t}$. By Proposition 2.4, there is a positive integer $M = Dp^r$ such that for any finite place v ,

$$v(t) < -v(M) \implies I_v(x, f) = 0.$$

In fact that Proposition says that for $v = N$, $I_v(x, f) \neq 0 \implies N \mid (t-1)$. Also, $I_\infty(x, f) = 0$ unless $x > 0$. Putting these together, we see that $I(x, f)$ is zero unless Mt is an integer $\neq 0, 1$, which is divisible by N . Thus

$$I_{\text{reg}}(f) := \sum_{n \neq 0, 1, N \mid (n-M)} I\left(\frac{n-M}{n}, f\right).$$

Lemma 7. *For any $n \neq 0$,*

$$I\left(\frac{n-M}{n}, f\right) << \frac{1}{n^{k/2-\varepsilon}},$$

with the implied constant independent of n ,

Proof. By Prop. 2.4, for any (finite) prime q , $I_q(\frac{n-M}{n}, f)$ is 1 for $q \nmid n(n-M)$, and if $q \mid n(n-M)$,

$$I_q\left(\frac{n-M}{n}, f\right) \leq Mv_q((n-M)/n)^2.$$

We *claim* that the function

$$g(n) := \prod_q v_q(n)$$

satisfies, for every $\varepsilon > 0$, the bound

$$g(n) \leq Cn^\varepsilon,$$

for a constant $C > 0$ independent of n .

To see this, first note that $g(n)$ is multiplicative in n (though not strictly). Fix any $\varepsilon > 0$. Then $\exists A > 0$ such that for any prime q and positive integer $a > 0$,

$$a \leq p^{a\varepsilon} \quad \text{if either } a > M \quad \text{or} \quad p > M.$$

Given $n > 0$ with unique factorization $n = \prod_{j \in J} q_j^{a_j}$, where the q_j are primes and $a_j > 0$, we may write $n = mk$, where m has no prime divisors $q_j < A$ with exponents $a_j < A$. It follows immediately that $g(m) \leq nm^\varepsilon$, and k is a product over primes $\leq A$ with exponents at most A . We obtain

$$g(n) = g(m)g(k) \leq Cn^\varepsilon$$

where C is the maximum of $g(k)$ for the finitely many choices of k .

Hence the *claim*.

It follows that for any $\varepsilon > 0$,

$$\prod_{q|n(n-M)} v_q((n-M)/n)^2 \ll n^\varepsilon.$$

Moreover, thanks to Prop. 2.4, we have

$$I_\infty\left(\frac{n-M}{n}, f\right) \ll \frac{1}{n^{k/2}}.$$

In both estimates, the implied constants depends on M , but *not* on N . This proves the Lemma. □

Consequently,

$$I_{\text{reg}}(f) \ll \sum_{n \neq 0, 1, N|n-M} \frac{1}{n^{k/2-\varepsilon}}.$$

But for any real $u > 1$,

$$\sum_{n \neq 0, 1, N|n-M} \frac{1}{n^u} = \sum_{m \geq 1} \frac{1}{(mN + M)^u} \ll \frac{1}{N^u}.$$

The Proposition now follows. □

4. The spectral side

Let $I(f)$ be the limit, as $s_1, s_2 \rightarrow 0$, of the integral of the kernel of the test function over the square of the torus $\tilde{T}$ against the quasi-character $(\chi|\cdot|^{s_1}, |\cdot|^{s_2})$ with χ odd. Let ρ denote the right regular representation of $\tilde{G}(\mathbb{A})$ on the space of cusp forms with trivial central character.

Let $\{\varphi_j\}$ denote an orthogonal basis of cusp forms of level N and weight $k \geq 2$. They can be chosen to be Hecke eigenforms, i.e., generate an admissible representation $\pi_j = \pi_{j,\infty} \otimes \pi_{j,0}$ of $\mathrm{GL}(2, \mathbb{A})$ satisfying $\pi_{j,\infty} \simeq \mathcal{D}_k$ and $\pi_{j,0}^{K_0(N)} \neq 0$. For the subspace of newforms of level N , we may take φ_j to be newforms. If $k < 12$, then we have only newforms.

Write $f = f_\infty \otimes f_0$, where f_0 denotes the finite part of f . By construction f_∞ is $d(\mathcal{D}_k)$ times the complex conjugate of the matrix coefficient h_∞ , say, of $\mathcal{D}$. When we replace the Haar measure dg by cdg , the formal degree gets multiplied by c^{-1} . The matrix coefficient h_∞ is independent of the choice of Haar measure, but the operator $\rho_\infty(h_\infty)$ does depend on the Haar measure. By contrast, the function f_∞ depends on the Haar measure, but the operator $\rho_\infty(f_\infty)$ is independent of the Haar measure. In fact, $\rho_\infty(f_\infty)$ is a projection operator onto the subspace of cusp forms of weight k . Note also that the kernel of a fixed function $h(g)$ does not depend on dg since it's just the sum $\sum h(g^{-1}\gamma h)$. However, the kernel for $\rho(f)$ depends inversely on Haar measure because the formal degree does.

Putting these remarks together, we obtain the spectral expression

$$K_f(x, y) = \sum_j \frac{\rho(f_0)\varphi_j(x)\overline{\varphi_j(y)}}{\langle \varphi_j, \varphi_j \rangle},$$

where the sum is over an orthogonal basis of cusp forms of weight k and level N . On each φ_j , $\rho(f)$ acts by $\pi_j(f)$. There is an adelic scalar product $\langle \cdot, \cdot \rangle_{\mathbb{A}}$ on the space of cusp forms, given by

$$\langle \varphi, \psi \rangle_{\mathbb{A}} = \int_{Z(\mathbb{A})G(\mathbb{Q}) \backslash G(\mathbb{A})} \varphi(g)\overline{\psi(g)}dg,$$

where dg denotes the quotient measure on $Z(\mathbb{A})G(\mathbb{Q}) \backslash G(\mathbb{A})$ induced by the Haar measure on $G(\mathbb{A})$, normalized so that the quotient space gets volume 1. One has the identification

$$Z(\mathbb{A})G(\mathbb{Q}) \backslash G(\mathbb{A}) / K_\infty K_0(N) = \Gamma_0(N) \backslash \mathcal{H},$$

and if φ is a cusp form of level N , $|\varphi|^2$ is right invariant by $K_\infty K_0(N)$. Consequently, $\langle \varphi, \varphi \rangle_{\mathbb{A}}$ is V_N times the Petersson scalar product $\langle \varphi, \psi \rangle$ on the classical

modular forms of level N . By abuse of notation, we are using the same symbol for the adelic cusp form and the classical modular form it defines.

Proposition 4.1. *We have*

$$I(f) = \frac{1}{4\pi V_N} \sum_j \frac{L(1/2, \varphi_j) L(1/2, \varphi_j \otimes \chi)}{\langle \varphi_j, \varphi_j \rangle} \hat{f}_p(\varphi_j),$$

where $\hat{f}_p(\varphi_j)$ denotes a local expression depending only on f_p and φ_j . For a suitable local function $f_{p,0}$,

$$\hat{f}_{p,0}(\varphi_j) = a_p(\varphi_j).$$

Proof. . Using the expression above for the kernel in conjunction with the definition of $I(f)$, we obtain

$$I(f) = \lim_{s_1 \rightarrow 0, s_2 \rightarrow 0} \sum_j P_j(f, \chi, s_1) Q_j(f, s_2),$$

where P_j, Q_j are period integrals given by

$$P_j(f, \chi, s_1) = \int_{\mathbb{Q}^* \backslash \mathbb{A}^*} \rho(f) \varphi_j \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \chi(a) |a|^{s_1} d^* a$$

and

$$Q_j(f, s_2) = \int_{\mathbb{Q}^* \backslash \mathbb{A}^*} \overline{\varphi}_j \begin{pmatrix} b & 0 \\ 0 & 1 \end{pmatrix} |a|^{s_2} d^* b.$$

If W_j denotes the Whittaker function of φ_j , there is a well known Fourier expansion (for $g \in \mathrm{GL}(2, \mathbb{A})$)

$$\varphi_j(g) = \sum_{t \in \mathbb{Q}^*} W_j \left[\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} g \right].$$

Since χ and the adelic absolute value $|\cdot|$ are 1 on $\mathbb{Q}^*$, we can unfold the expression for P_j as

$$P_j(f, \chi, s_1) = \int_{\mathbb{A}^*} \rho(f) W_j \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \chi(a) |a|^{s_1} d^* a$$

Then from the factorizability of W_j , f and χ , we get the product expansion

$$P_j(f, \chi, s_1) = \prod_v P_{j,v}(f, \chi, s_1),$$

where v runs over all the places of $\mathbb{Q}$, with

$$P_{j,v}(f, \chi, s_1) = \int_{\mathbb{Q}_v^*} \rho(f_v) W_{j,v} \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \chi_v(a) |a|^{s_1} d^* a.$$

For $v \neq p$, f_v is chosen such that $\rho(f_v)W_{j,v}$ times $\chi_v \circ \det$ is the new vector of $\pi_{j,v} \otimes \chi_v$, so that by Jacquet-Langlands,

$$P_{j,v}(f, \chi, s_1) = L(s_1 + 1/2, \pi_{j,v} \otimes \chi_v).$$

This is clear at any v where χ is unramified. Let us indicate the reason at a $v = q$ where χ is ramified, say of conductor q^m . It follows from the definition of f_v , and the transformation property of the Whittaker function under the left translation by the maximal unipotent subgroup, that

$$P_{j,v}(f, \chi, s_1) = \frac{1}{g(\chi_v)} \sum_{n \in \mathbb{Z}} W_v \begin{pmatrix} \varpi^n & 0 \\ 0 & 1 \end{pmatrix} g(\chi_v, \psi_{\varpi^{n-m}}) q^{-n(s_1+1/2)},$$

where $\psi_t(x) = \psi(tx)$ and ψ the additive character defined as the composite

$$\mathbb{Q}_q \rightarrow \mathbb{Q}_q/\mathbb{Z}_q \rightarrow \mathbb{Q}/\mathbb{Z} \rightarrow S^1,$$

with the last arrow on the right being $x \rightarrow e^{2\pi i x}$. Note that $g(\chi_v)$ is just $g(\chi_v, \psi_{\varpi^{-m}})$. Since W_v has been chosen to be the spherical vector giving the right L -factor of π_v , we have

$$W_v \begin{pmatrix} \varpi^n & 0 \\ 0 & 1 \end{pmatrix} = \delta_{n \geq 0} q^{n/2} \left(\frac{\alpha^{n+1} - \beta^{n+1}}{\alpha - \beta} \right),$$

where $\delta_{n \geq 0}$ is 1 if $n \geq 0$ and is zero otherwise, and $\text{diag}\{\alpha, \beta\}$ is the conjugacy class attached to π_v . (In other words, $L(s, \pi_v)$ is the inverse of $(1 - \alpha q^{-s})(1 - \beta q^{-s})$.) So we may take n to be non-negative from here on. On the other hand, it is well known that $g(\chi_v, \psi_{\varpi^r}) = 0$ is $r < m$. Taking $r = m - n$ we get vanishing for $n > 0$. Thus $n = 0$, and we get (for any s_1)

$$P_{j,v}(f, \chi, s_1) = 1,$$

To finish, note that

$$L(s, \pi_v \otimes \chi_v) = 1,$$

because π_v is unramified, while χ_v is ramified.

Similarly, $\forall v \neq p$,

$$Q_j(f, s_2) = \prod_v Q_{j,v}(f, s_2),$$

with

$$Q_{j,v}(f, s_2) = L(\bar{s}_2 + 1/2, \pi_{j,v}), \forall v.$$

The main assertion of the Proposition now follows, after letting s_1 and s_2 go to 0, which is justified because the expressions are known to have analytic continuations, with

$$\hat{f}_p(\varphi_j) = \frac{P_{j,v}(f, \chi, 0)}{L(1/2, \pi_{j,p} \otimes \chi_p)}.$$

Finally, one knows that there is a Hecke function $f_{p,0}$ such that

$$\pi_j(f_{p,0})\varphi_j = a_p(\pi_j)\varphi_j,$$

and this gives the final assertion. □

5. Proof of the Main Theorem sans measure

Now we will exploit the equality of the geometric and spectral sides of the relative trace formula at hand, for forms of even weight $k > 2$ and prime level N . Continue to assume that χ is a quadratic character of discriminant $D < 0$.

The sum of the regular terms on the geometric side was estimated in section 3. There are six singular terms, and four of them vanish. As $s_j \rightarrow 0$, for $j = 1, 2$, each of the remaining pair of singular terms, which are the dominant ones, goes to $L(1, \chi)$. In view of the expression for the spectral side (cf. section 4), we then obtain the following (for any $\varepsilon > 0$):

$$\frac{1}{4\pi} \sum_{\varphi \in \mathcal{F}_{N,k}} \frac{1}{(\varphi, \varphi)} L\left(\frac{1}{2}, \varphi \otimes \chi\right) L\left(\frac{1}{2}, \varphi\right) = 2c_k L(1, \chi) + O(N^{-k/2+\varepsilon}),$$

where $\mathcal{F}_{N,k}$ is an orthogonal basis of holomorphic, Hecke eigencusp forms of weight k and level N , and c_k is as in the statement of Theorem A.

Now let $\mathcal{F}_N(k)^{\text{new}}$ be the subspace of newforms of level N and even weight k . The sum on the left of the formula above runs over newforms of level N and oldforms which necessarily come from level 1. Since $\chi(-1) = -1$, we see that the sign of the functional equation of $L(s, g \otimes \chi)L(s, g)$ is -1 for any form of level 1, implying that $L(1/2, g \otimes \chi)L(1/2, g)$ is zero for any such g . There are two oldforms of level N associated to g , which, in classical language, are $g(z)$ and $g_N(z) := g(Nz)$. It is immediate that their Mellin transforms are related by $L(s, g_N) = N^{1-s}L(s, g)$. It follows that $L(1/2, g_N \otimes \chi)L(1/2, g_N)$ is also 0.

Now we have proved Theorem A without the assertion about the measure.

6. The measure

Fix $v = p$ not dividing ND , so that π and χ are both unramified at p .

Let f_n be the characteristic function of $ZK \begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix} K$ for $n = 0, 1, 2, \dots$ and let $\varphi_n(s)$ be the Satake transform of f_n . Then $\varphi_n(s)$ is a symmetric Laurent polynomial in p^s and p^{-s} with the property:

$$\text{Trace}(\pi_s(f_n)) = \varphi_n(s)$$

where π_s is the principal series representation unitarily induced from the character

$$\begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix} \rightarrow \left| \frac{\alpha}{\beta} \right|^s$$

Lemma 8. $\varphi_0(s) = 1$ and for $n \geq 1$,

$$\varphi_n(s) = p^{n/2} \left[p^{ns} + p^{-ns} + \left(1 - \frac{1}{p} \right) \left(p^{(n-2)s} + p^{(n-4)s} + \dots + p^{-(n-4)s} + p^{-(n-2)s} \right) \right]$$

Proof. The coset $K \begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix} K$ is equal to a union of single cosets gK where g ranges over the following elements:

$$\begin{aligned} \begin{pmatrix} p^n & t \\ 0 & 1 \end{pmatrix} & \quad t \in \mathcal{O}/(p^n) \\ \begin{pmatrix} p^{n-k} & t \\ 0 & p^k \end{pmatrix} & \quad t \in \left(\mathcal{O}/p^{n-k} \right)^* \quad k = 1, \dots, n \end{aligned}$$

Using this we easily calculate the action of f_n on the principal series π_s and we find the above formula. $\square$

We want to find the Satake transform of the following linear functional for bi- K -invariant f :

$$T(f) = I_v(n^+) = \iint_{\mathbb{Q}_v^* \times \mathbb{Q}_v^*} f\left(\begin{pmatrix} b & a \\ 0 & 1 \end{pmatrix}\right) \chi_v |a|^{-s_1-s_2} \chi_{2,v}(b) |b|^{s_2} d^*a d^*b.$$

In fact, we will take χ_2 trivial, $s_2 = 0$, and set $s = s_1$. We assume χ is unramified, so we have:

$$T(f) = I_v(n^+) = \iint_{\mathbb{Q}_v^* \times \mathbb{Q}_v^*} f\left(\begin{pmatrix} b & a \\ 0 & 1 \end{pmatrix}\right) \chi(a) |a|^{-s} d^*a d^*b.$$

We first evaluate $T(f_n)$ for $n \geq 0$. We first note that if

$$\begin{pmatrix} b & a \\ 0 & 1 \end{pmatrix} \in ZK \begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix} K$$

then

$$\begin{pmatrix} b & a \\ 0 & 1 \end{pmatrix} \in K \begin{pmatrix} p^{n+\alpha} & 0 \\ 0 & p^\alpha \end{pmatrix} K$$

for some α . This is not possible unless $\alpha \leq 0$ and $b = p^{n+2\alpha}u$ for some unit u . Furthermore, the gcd of $\{a, p^{n+2\alpha}, 1\}$ is p^α , so we must have $n+2\alpha \geq \alpha$, that is, $\alpha \geq -n$.

If $0 > \alpha > -n$, then the gcd condition forces $a \in p^\alpha \mathbb{Z}_p^*$. If $\alpha = 0$, then we just have $a \in \mathbb{Z}_p^*$ and if $\alpha = -n$, we have $a \in p^{-n} \mathbb{Z}_p^*$.

Set $\delta = \chi(p)$. For $n = 0$, we have:

$$T(f_0) = \int_{u \in \mathbb{Z}_p^*} \int_{a \in \mathbb{Z}_p} \chi(a) |a|^{-s} d^*u d^*a = L_p(-s, \chi)$$

For $n > 0$, $T(f_n)$ is a sum of three terms:

$$I = \int_{u \in \mathbb{Z}_p^*} \int_{a \in \mathbb{Z}_p} f\left(\begin{pmatrix} p^n u & a \\ 0 & 1 \end{pmatrix}\right) \chi(a) |a|^{-s} d^*u d^*a = L_p(-s, \chi)$$

and

$$II = \int_{u \in \mathbb{Z}_p^*} \int_{a \in p^{-n}\mathbb{Z}_p} f\left(\begin{pmatrix} p^{-n}u & a \\ 0 & 1 \end{pmatrix}\right) \chi(a) |a|^{-s} d^*u d^*a = \delta^{-n} p^{-ns} L_p(-s, \chi)$$

$$\begin{aligned} III &= \sum_{\alpha=1}^{n-1} \int_{u \in \mathbb{Z}_p^*} \int_{v \in \mathbb{Z}_p^*} f\left(\begin{pmatrix} p^{n-2\alpha}u & p^{-\alpha}v \\ 0 & 1 \end{pmatrix}\right) \chi(p^{-\alpha}) |p^{-\alpha}|^{-s} d^*u d^*v \\ &= \sum_{\alpha=1}^{n-1} \chi(p^{-\alpha}) p^{-\alpha s} = \frac{\delta^{-1} p^{-s} - \delta^{-n} p^{-ns}}{1 - \delta^{-1} p^{-s}} = \frac{\delta^{1-n} p^{(1-n)s} - 1}{1 - \delta p^s} \\ &= L_p(-s, \chi) (\delta^{1-n} p^{(1-n)s} - 1) \end{aligned}$$

So we get the sum:

$$T(f_n) = L_p(-s, \chi) (\delta^{-n} p^{-ns} + \delta^{1-n} p^{(1-n)s}) = \delta^{-n} p^{-ns} L_p(-s, \chi) (1 + \delta p^s)$$

Note that if $\delta = -1$, then $T(f_n)$ this approaches 0 as $s \rightarrow 0$ and in this, the limit is the *Plancherel measure* at p , that is, the measure μ_- satisfying $\mu(f_0) = 1$ and $\mu(f_n) = 0$ for $n > 0$.

Now we calculate

$$S(f) = I_v(n^-) = \iint_{\mathbb{Q}_v^* \times \mathbb{Q}_v^*} f\left(\begin{pmatrix} a & 0 \\ b & 1 \end{pmatrix}\right) \chi(a^{-1}b) |a|^{-s_1} |b|^{s_1+s_2} d^*a d^*b.$$

We evaluate $S(f_n)$ for $n \geq 0$. We first note that if

$$\begin{pmatrix} a & 0 \\ b & 1 \end{pmatrix} \in ZK \begin{pmatrix} p^n & 0 \\ 0 & 1 \end{pmatrix} K$$

then

$$\begin{pmatrix} a & 0 \\ b & 1 \end{pmatrix} = \begin{pmatrix} p^n u & 0 \\ b & 1 \end{pmatrix}, \begin{pmatrix} p^{n-2\alpha} u & 0 \\ p^{-\alpha} v & 1 \end{pmatrix} \text{ or } \begin{pmatrix} p^{-n} u & 0 \\ p^{-n} b & 1 \end{pmatrix}$$

where $b \in \mathbb{Z}_p$ and u, v are units in $\mathbb{Z}_p^*$, and $0 \leq \alpha \leq n$.

For $n = 0$, we have:

$$T(f_0) = \int_{u \in \mathbb{Z}_p^*} \int_{b \in \mathbb{Z}_p} \chi(b) |b|^{s_1+s_2} d^*u d^*a = L_p(s_1 + s_2, \chi)$$

For $n > 0$, $T(f_n)$ is a sum of three terms:

$$\begin{aligned}
 I &= \int_{u \in \mathbb{Z}_p^*} \int_{b \in \mathbb{Z}_p} \chi(p^{-n}b) |p^n|^{-s_1} |b|^{s_1+s_2} d^*u d^*b = \delta^{-n} p^{ns_1} L(s_1 + s_2, \chi) \\
 II &= \sum_{\alpha=1}^{n-1} \int_{u \in \mathbb{Z}_p^*} \int_{v \in \mathbb{Z}_p^*} \chi(p^{\alpha-n}) |p^{n-2\alpha}|^{-s_1} |p^{-\alpha}|^{s_1+s_2} d^*u d^*v \\
 &= p^{ns_1} \delta^{-n} \sum_{\alpha=1}^{n-1} \delta^\alpha p^{\alpha(s_2-s_1)} = p^{ns_1} \delta^{-n} \frac{\delta p^{s_2-s_1} - \delta^2 p^{n(s_2-s_1)}}{1 - \delta p^{s_2-s_1}} = p^{ns_1} \delta^{-n} \frac{\delta p^{s_2-s_1} - \delta^2 p^{s_2}}{1 - \delta p^{s_2}} \\
 III &= \int_{u \in \mathbb{Z}_p^*} \int_{b \in \mathbb{Z}_p} \chi(b) |p^{-n}|^{-s_1} |p^{-n}b|^{s_1+s_2} d^*a d^*b = p^{ns_2} L(s_1 + s_2, \chi)
 \end{aligned}$$

When we go from $I_v(n^+)$ to $I_v(n^-)$, the main change is that (s_1, s_2) gets sent to $(-s_2, -s_1)$. We get $I_v(n^+) = I_v(n^-)$ (as $s_1, s_2 \rightarrow 0$), and we again get the Plancherel measure at p when $\chi(p) = -1$.

We are left to analyze the case $\chi(p) = 1$. We will simply write μ for μ_+ . By the calculation above, we have

$$T(f_0) = 1, \quad \text{and} \quad T(f_n) = \delta^n + \delta^{-n} \quad \forall n > 0,$$

with $\delta = 1$. It will be convenient (and more transparent) for us to carry out the calculations for arbitrary non-zero scalar δ and at the end set $\delta = 1$. We seek a function $F(s)$ of p^s such that

$$\int \varphi_n(s) F(s) ds = \begin{cases} 1 & n = 0 \\ \delta^n + \delta^{-n} & n \geq 1 \end{cases}$$

The integral is taken over $s \in i\mathbf{R}/(2\pi i \ln p \mathbf{Z})$.

Proposition 6.1.

$$F_{ev}(s) = 1 + \sum_{n=1}^{\infty} A_n p^{2ns-n}$$

where

$$\begin{aligned}
 A_n &= \delta^{2n} + \delta^{-2n} - (p-1) \left(\delta^{2n-2} + \delta^{2n-4} + \delta^{2n-6} + \dots + \delta^{-2n+4} + \delta^{-2n+2} \right) \\
 &= \delta^{2n} + \delta^{-2n} - (p-1) \left(\frac{\delta^{2n-1} - \delta^{-2n+1}}{\delta - \delta^{-1}} \right)
 \end{aligned}$$

Proof. Observe that

$$\int p^{ns} p^{ms} ds = \begin{cases} 1 & n + m = 0 \\ 0 & \text{otherwise} \end{cases}$$

Therefore $\int \varphi_{2n}(s) F_{ev}(s) ds$ is equal to

$$A_n + \left(1 - \frac{1}{p}\right) \left[pA_{n-1} + p^2A_{n-2} + \cdots p^n A_0 \right] = \delta^{2n} + \delta^{-2n} + \sum_{j=-n+1}^{n-1} c_j \delta^{2j}$$

We must show that the coefficients c_j are zero. We have:

$$\begin{aligned} c_j &= -(p-1) - \left(1 - \frac{1}{p}\right) \left[p(p-1) + p^2(p-1) + \cdots + p^{j-1}(p-1) - p^j \right] \\ &= -(p-1) - \left(1 - \frac{1}{p}\right) (-p) = 0 \end{aligned}$$

□

Proposition 6.2.

$$F_{od}(s) = B_0 p^{s-\frac{1}{2}} + p^{3s-\frac{3}{2}} B_1 + p^{5s-\frac{5}{2}} B_2 + \cdots = \sum_{n \geq 1} B_n p^{(2n+1)s-(n+\frac{1}{2})}$$

where

$$\begin{aligned} B_n &= \delta^{2n+1} + \delta^{-2n-1} - (p-1) \left(\delta^{2n-1} + \delta^{2n-3} + \delta^{2n-5} + \cdots + \delta^{-2n+3} + \delta^{-2n+1} \right) \\ &= \delta^{2n+1} + \delta^{-2n-1} - (p-1) \left(\frac{\delta^{2n} - \delta^{-2n}}{\delta - \delta^{-1}} \right) \end{aligned}$$

Proof. A similar computation gives that $\int \varphi_{2n+1}(s) F_{ev}(s) ds$ is equal to

$$B_n + \left(1 - \frac{1}{p}\right) \left[pB_{n-1} + p^2B_{n-2} + \cdots p^n B_0 \right] = \delta^{2n+1} + \delta^{-2n-1} + \sum_{j=-n+1}^{n-1} c_j \delta^{2j+1}$$

□

□

In short, we get

$$F(s) = 1 + \sum_{n=1}^{\infty} C_n p^{n(s-\frac{1}{2})}$$

where

$$C_n = \delta^n + \delta^{-n} - (p-1) \left(\frac{\delta^{n-1} - \delta^{-n+1}}{\delta - \delta^{-1}} \right)$$

To write $F(s)$ as a rational function, we sum the geometric series. Let $T = p^{(s-\frac{1}{2})}$.

$$\begin{aligned}
F(s) &= 1 + \sum_{n \geq 1} \left[(\delta T)^n + (\delta^{-1} T)^n - \frac{(p-1)}{\delta - \delta^{-1}} \left(\delta^{-1} (\delta T)^n + \delta (\delta^{-1} T)^n \right) \right] \\
&= 1 + \sum_{n \geq 1} \frac{(\delta T)^n}{\delta - \delta^{-1}} \left[\delta - \delta^{-1} - p\delta^{-1} + \delta^{-1} \right] + \sum_{n \geq 1} \frac{\delta^{-1} T^n}{\delta - \delta^{-1}} \left[\delta - \delta^{-1} + p\delta - \delta \right] \\
&= 1 + \frac{\delta - p\delta^{-1}}{\delta - \delta^{-1}} \sum_{n \geq 1} (\delta T)^n - \frac{\delta^{-1} - p\delta}{\delta - \delta^{-1}} \sum_{n \geq 1} (\delta^{-1} T)^n \\
&= 1 + \frac{\delta - p\delta^{-1}}{\delta - \delta^{-1}} \frac{\delta T}{1 - \delta T} - \frac{\delta^{-1} - p\delta}{\delta - \delta^{-1}} \frac{\delta^{-1} T}{1 - \delta^{-1} T}
\end{aligned}$$

Using Mathematica, we find the further simplification:

$$F(s) = \frac{1 - pT^2}{(T - \delta)(T - \delta^{-1})} = \frac{1 - p^{2s}}{(1 - \delta^{-1} p^{s-\frac{1}{2}})(1 - \delta p^{s-\frac{1}{2}})}$$


We can also replace $F(s)$ by $\Re(F(s)) = \frac{1}{2}(F(s) + F(-s))$ (for s purely imaginary). We find that

$$\mu(s) = \frac{1}{2}(F(s) + F(-s)) = \frac{1}{2} \frac{\left(1 - \frac{1}{p}\right)(2 - p^{2s} - p^{-2s})}{\left(1 + \frac{1}{p} - p^{-\frac{1}{2}+s} - p^{-\frac{1}{2}-s}\right)^2}$$

The linear functional is

$$T(f) = \frac{1}{T} \int_0^T \widehat{f}(is) \mu(is) ds$$

where $T = \pi / \log(p)$. Here is a graph of $\mu(s)$ for the case $p = 2$ and δ trivial.



Now let $x = p^s + p^{-s} = 2 \cos(s \log p)$. Then $x^2 = 2 + p^{2s} + p^{-2s}$ and we can also write

$$\mu(s) = \frac{1}{2}(F(s) + F(-s)) = \frac{1}{2} \frac{\left(1 - \frac{1}{p}\right)(4 - x^2)}{\left(1 + \frac{1}{p} - p^{-\frac{1}{2}}x\right)^2}$$

$$dx = -2 \log(p) \sin(s \log p) ds = -2 \log(p) \sqrt{1 - (x/4)^2} ds = -\log(p) \sqrt{4 - x^2} ds$$

and thus

$$\begin{aligned} \frac{\log(p)}{\pi} \mu(s) ds &= - \left(\frac{\log(p)}{\pi} \right) \frac{1}{2 \log(p)} \frac{\left(1 - \frac{1}{p}\right) \sqrt{4 - x^2}}{\left(1 + \frac{1}{p} - p^{-\frac{1}{2}}x\right)^2} dx \\ &= - \frac{\pi}{2} \frac{\left(1 - \frac{1}{p}\right) \sqrt{4 - x^2}}{\left(1 + \frac{1}{p} - p^{-\frac{1}{2}}x\right)^2} dx \end{aligned}$$

Notice that the minus sign disappears when we write the integral as an integral over $[-2, 2]$.

REFERENCES

- [A-S] M. Abramowitz and I.A. Stegun, *Handbook of mathematical functions*, Dover edition (1965).
- [Du] W. Duke, The critical order of vanishing of automorphic L -functions with large level, *Inventiones Math.* **119**, 165-174 (1995).
- [GJ] S. Gelbart and H. Jacquet, A relation between automorphic representations of $GL(2)$ and $GL(3)$, *Ann. Scient. Éc. Norm. Sup.* (4) **11** (1979), 471-542.
- [Gu] J. Guo, On the positivity of the central critical values for automorphic L -functions for $GL(2)$, *Duke Mathematical Journal* **83**, No. 1, 157-190 (1996).
- [IwS] H. Iwaniec and P. Sarnak, The non-vanishing of central values of automorphic L -functions and Landau-Siegel zeros, *Israel Journal of Math* **120**, 155-177 (2000).
- [J] H. Jacquet, Sur une résultat de Waldspurger, *Ann. Scient. Éc. Norm. Sup.*, 4^e série, t. **19**, 185-229 (1986).
- [KMV] E. Kowalski, P. Michel, J. VanderKam, Non-vanishing of high derivatives of automorphic L -functions at the center of the critical strip, *J. Reine Angew. Math.* **526** 1-34 (2000).
- [Lu] W. Luo, On the non-vanishing of Rankin-Selberg L -functions, *Duke Math Journal* **69**, 411-427 (1993).
- [Sa] P. Sarnak, Statistical properties of eigenvalues of the Hecke operators, in *Analytic number theory and Diophantine problems*, 321-331, *Progr. Math.* **70**, Birkhuser, Boston (1987).
- [Se] J.-P. Serre, Répartition asymptotique des valeurs propres de l'opérateur de Hecke T_p , *Journal of the AMS* **10**, no. 1, 75-102 (1997).

Dinakar Ramakrishnan

Professor of Mathematics, 253-37 Caltech, Pasadena, CA 91125, USA

E-mail: dinakar@its.caltech.edu

Jonathan Rogawski

Professor of Mathematics Department of Mathematics

University of California, Los Angeles, California 90095

E-mail: jonr@math.ucla.edu